

The Araucaria Project. The Distance to the Local Group Galaxy WLM from Near-Infrared Photometry of Cepheid Variables ¹

Wolfgang Gieren

*Universidad de Concepción, Departamento de Física, Astronomy Group, Casilla 160-C,
Concepción, Chile*

wgieren@astro-udec.cl

Grzegorz Pietrzyński

*Universidad de Concepción, Departamento de Física, Astronomy Group, Casilla 160-C,
Concepción, Chile*

Warsaw University Observatory, Al. Ujazdowskie 4, 00-478, Warsaw, Poland

pietrzyn@astrouw.edu.pl

Olaf Szewczyk

*Universidad de Concepción, Departamento de Física, Astronomy Group, Casilla 160-C,
Concepción, Chile*

szewczyk@astro-udec.cl

Igor Soszyński

Warsaw University Observatory, Al. Ujazdowskie 4, 00-478, Warsaw, Poland

soszynsk@astrouw.edu.pl

Fabio Bresolin

*Institute for Astronomy, University of Hawaii at Manoa, 2680 Woodlawn Drive, Honolulu
HI 96822, USA*

bresolin@ifa.hawaii.edu

Rolf-Peter Kudritzki

*Institute for Astronomy, University of Hawaii at Manoa, 2680 Woodlawn Drive, Honolulu
HI 96822, USA*

kud@ifa.hawaii.edu

Miguel A. Urbaneja

*Institute for Astronomy, University of Hawaii at Manoa, 2680 Woodlawn Drive, Honolulu
HI 96822, USA*

`urbaneja@ifh.hawaii.edu`

Jesper Storm

Astrophysikalisches Institut Potsdam, An der Sternwarte 16, D-14482 Potsdam, Germany

`jstorm@aip.de`

Dante Minniti

*Departamento de Astronomia y Astrofisica, Pontificia Universidad Católica de Chile,
Casilla 306, Santiago 22, Chile*

`dante@astro.puc.cl`

ABSTRACT

We have obtained deep images in the near-infrared J and K filters for several fields in the Local Group galaxy WLM. We report intensity mean magnitudes for 31 Cepheids located in these fields which we previously discovered in a wide-field optical imaging survey of WLM. The data define tight period-luminosity relations in both near-infrared bands which we use to derive the total reddening of the Cepheids in WLM and the true distance modulus of the galaxy from a multiwavelength analysis of the reddened distance moduli in the VIJK bands. From this, we obtain the values $E(B-V) = 0.082 \pm 0.02$, and $(m-M)_0 = 24.924 \pm 0.042$ mag, with a systematic uncertainty in the distance of about $\pm 3\%$. This Cepheid distance agrees extremely well with the distance of WLM determined from the I-band TRGB method by ourselves and others. Most of the reddening of the Cepheids in WLM (0.06 mag) is produced inside the galaxy, demonstrating again the need for an accurate determination of the total reddening and/or the use of infrared photometry to derive Cepheid distances which are accurate to 3% or better, even for small irregular galaxies like WLM.

Subject headings: distance scale - galaxies: distances and redshifts - galaxies: individual(WLM) - stars: Cepheids - infrared photometry

1. Introduction

The effectiveness of using multiwavelength optical and near-infrared (NIR) observations of Cepheid variables for distance determination of galaxies has been known for a long time (McGonegal et al. 1982; Madore & Freedman 1991). Only recently however the technical problems with obtaining accurate and reliable NIR photometry for faint objects in dense regions have been solved. Using NIR observations of Cepheids provides a number of advantages for accurate distance work. First, the total and differential reddening is significantly reduced in comparison to the optical bandpasses. Second, the Cepheid period-luminosity (PL) relation becomes steeper toward longer wavelengths, and its intrinsic dispersion becomes smaller (e.g. Fouqué et al. 2007), both factors helping in deriving a more accurate distance. Third, metallicity effects on the PL relation in the near-IR are expected to be less important than at optical wavelengths (Bono et al. 1999). Fourth, and very importantly from an observational point of view, the amplitudes of variability are significantly smaller in the NIR than at optical wavelengths, so even random single-epoch observations approximate the mean magnitude reasonably well. If the period and optical light curve of a Cepheid is accurately known, it is possible to derive its mean magnitude in the NIR bands with an impressive 1-2% accuracy from just one single-epoch observation (Soszyński et al. 2005).

Simultaneously studying NIR and optical PL relations of Cepheids provides another important advantage. By combining the reddened distance moduli in the optical and NIR it is possible to derive both the total reddening and the true distance modulus with very good accuracy. This has been demonstrated in the previous papers of our ongoing Araucaria Project which reported the distances to NGC 300 (Gieren et al. 2005b), IC 1613 (Pietrzyński et al. 2006), NGC 6822 (Gieren et al. 2006), NGC 3109 (Soszyński et al. 2006) and NGC 55 (Gieren et al. 2008) derived by this method. For all these galaxies, we were able to determine their distances *with respect to the LMC* with a 3-5% accuracy from our technique. The possibility to measure the reddening produced *inside* these galaxies has been a fundamental step towards achieving this accuracy. We recall here that the principal purpose of the Araucaria Project is to measure the distances to a sample of nearby galaxies with widely different environmental properties with different stellar distance indicators, and compare the results to determine the metallicity and age dependences of each standard candle.

In a previous paper (Pietrzyński et al. 2007; hereafter Paper I), we reported on an extensive wide-field imaging survey for Cepheids in the WLM irregular galaxy carried out in the optical V and I bands in which we have discovered 60 variables, greatly extending the

¹Based on observations obtained with the ESO NTT for Programmes 077.D-0423 and 080.D-0047(B) and with the Magellan telescope at the Las Campanas Observatory

small sample of 15 Cepheids which had been previously discovered in WLM by Sandage & Carlson (1985). The WLM irregular galaxy is an important target of the Araucaria Project because of its low metallicity environment which allows to study the behavior of stellar distances indicators in the very low-metallicity regime. Indeed, the young stellar population of WLM with a mean metallicity of -0.8 dex (Bresolin et al. 2006; Urbaneja et al. 2008), is slightly more metal-poor than that of the SMC. From the VI optical photometry of the Cepheids we obtained in Paper I a true distance modulus of 25.144 mag for WLM. In the course of the work for the present paper, we discovered that unfortunately we had confused in Paper I the distance moduli for the V and Wesenheit bands; actually our adopted best value for the WLM true distance modulus from the reddening-free Wesenheit index in Paper I is 25.014 mag (and not 25.144 mag). In order to improve on the distance value derived from optical photometry of the Cepheids, it was clearly in order to extend the Cepheid distance work to the near-infrared domain and obtain a result which is unaffected by reddening. This is the strategy we have been employing in all the Araucaria work we have been conducting so far. In this paper, we therefore extend the light curve coverage for 31 Cepheids in WLM to the NIR J and K bands. We then utilize the multiband data available for these stars for an accurate determination of the total (average) interstellar extinction to the Cepheids in WLM, and determine an improved Cepheid distance to the galaxy from the multiwavelength method.

The paper is organized as follows. In section 2 we describe the NIR observations, data reductions and calibration of our photometry. In section 3 we derive the J- and K-band Cepheid PL relations in WLM from our data and determine the true distance modulus to the galaxy from a multiwavelength analysis. In section 4, we discuss our results, and in section 5 we summarize our conclusions.

2. Observations, Data Reduction and Calibration

The near-infrared data presented in this paper were collected during three observing runs in 2006 and 2007. We have been using two different telescopes and cameras for the data acquisition. On September 22 and 23, 2006, and on November 24 and 25, 2007 we used the SOFI infrared camera at the NTT at La Silla Observatory with a 4.9×4.9 arcmin field of view and a pixel scale of 0.288 arcsec/pixel, while on 22 and 23 November 2007 the PANIC infrared camera attached to the Baade Magellan Telescope at Las Campanas Observatory was used. For the PANIC observations, the field of view was 2.1×2.1 arcmin, and the pixel scale was 0.125 arcsec/pixel.

During the four nights at La Silla which were all photometric, we obtained deep J s and

*K*s observations of one field in WLM containing 23 Cepheid variables for which previous optical light curves were presented in Paper I. With the PANIC camera we were able to observe three additional smaller fields partly overlapping with the SOFI field, under photometric conditions. The locations of all these fields are shown in Fig. 1 and their coordinates are given in Table 1. Altogether, the four observed fields in WLM contain 31 Cepheids. In order to account for the rapid variations of the sky brightness in the near-infrared bands we used a dithering technique, as described in the previous papers of the Araucaria Project reporting infrared photometry of Cepheid variables. Total integration times were 15 min in the *J* and 60 min in the *K* band.

All the reductions and calibrations were performed with the pipeline developed in the course of the Araucaria Project and described in detail in earlier papers of this series. The subtraction of the sky brightness was done in a two-step process which included the masking of stars with the IRAF xdimsum package (Pietrzynski & Gieren 2002). Next, each single image was flat-fielded and stacked into the final deep field image. PSF photometry and aperture corrections were performed as described in Pietrzynski et al. (2002). The calibration of the photometry onto the standard system was based on the observations of 22 standard stars from the UKIRT list (Hawarden et al. 2001). All of them were observed along with the target fields in WLM at different airmasses and under photometric conditions. The large number of standard stars observed on each of the six photometric nights for this programme allowed us to obtain the absolute photometric zero points with an accuracy close to 0.01 mag, in both filters. Deviations of the photometric zero points obtained independently for different nights never exceeded 0.02 mag, for both SOFI and PANIC data. This is demonstrated in Fig. 2. For an external check of our photometry, we compared our calibrated magnitudes with those of the 2MASS catalog for the (small) sample of common stars, not finding any evidence for a significant zero point difference in *J* and/or *K* (see Fig. 3).

We present the calibrated individual near-infrared magnitudes for all Cepheids located in the observed fields in Table 2 which lists the stars’ IDs, heliocentric Julian days of the observations (mid-integrations), and the measurements in *J* and *K* with their standard deviations. Depending on the positions of the Cepheids in the galaxy, the number of individual JK observations range from one to six. On average, we were able to collect 3 JK observations per star. The WLM Cepheids observed in the near-infrared span a period range from 54 down to 2.5 days.

3. Near-Infrared Period-Luminosity Relations

The intensity mean magnitudes of the Cepheids were derived by taking a straight average of the individual random-phase fluxes measured of each variable, and converting the average fluxes back into magnitudes. In principle a more accurate procedure to determine the mean magnitudes in J and K would be the recipe described by Soszynski et al. (2005) which uses the V and I light curves of the variables and the known phases of the near-infrared observations. However, in the present case of WLM the epoch difference between the near-infrared data reported in this paper, and the previous VI data reported in Paper I is so large, typically some 200 pulsation cycles, that with the limited accuracy of the periods of the variables derived in Paper I the phasing of the near-IR data becomes very uncertain. While this is unfortunate, the simple taking of a straight average of several random-phase magnitudes in a near-IR band of a Cepheid still does produce a rather accurate mean magnitude, given the low light curve amplitudes of Cepheid variables at these wavelengths of typically 0.3 mag for stars with periods less than 10 days (e.g. Persson et al. 2004). For the one long-period Cepheid in WLM, cep001 with a period of 54 days, the amplitude of the K-band light curve is expected to be about 0.5 mag, but for this variable we have obtained six observations at different phases which makes us expect that their mean value is very close to the true mean magnitude of the variable, in the two bands we observed.

Table 3 gives the intensity mean J and K magnitudes of the individual Cepheids, with their estimated uncertainties from the number and accuracy of the individual observations leading to the adopted mean magnitude. We also provide the periods (adopted from Paper I). In Figures 4 and 5 we show the period-mean magnitude relations in the J and K bands as defined by the data in Table 3. There is one Cepheid, cep038, which is clearly over-luminous in both PL diagrams, by about 1.5 mag in J and about 2 mag in K. We assume that the very bright magnitude of this variable is caused by a nearby bright object which is not resolved in our images. Since the V and I magnitudes of cep038 are normal for its period (see Paper I), the blend must be a very red object. The occurrence of at least one strongly blended Cepheid in WLM in a sample of about 30 stars is quite expected from the result obtained by Bresolin et al. (2005) who studied the blending of Cepheids in NGC 300, at about twice the distance of WLM, from a comparison of ground-based and HST/ACS photometry, finding three strongly blended Cepheids in a sample of 16 Cepheids in this galaxy. For the following distance analysis, we exclude star cep038.

For the reasons discussed in Paper I, and in conformity with the approach adopted there, we adopt a period cutoff of $\log P$ (days) = 0.5 for the distance analysis, retaining only the Cepheids with longer periods in the sample. This ensures that possible overtone pulsators are likely to be excluded in the sample adopted for the distance determination, and

it eliminates the variables with the lowest signal-to-noise ratio in the photometry. The final sample consists of Cepheids 001-033 in Table 3 (24 stars). Note that the strongly blended Cepheid cep038 is eliminated from the final sample also on the basis of the adopted period cutoff.

From the 24 Cepheids in the final sample, least-squares fits to a line yield slopes of -3.067 ± 0.204 in K, and -3.073 ± 0.233 in J, respectively. These slope values are shallower than, but within 1σ consistent with the slopes for the Cepheid PL relations in the LMC, which are -3.261 in K, and -3.153 in J (Persson et al. 2004). Following the procedure we have used in our previous papers, we adopt the LMC slopes of Persson et al. (2004) in our fits. This yields the following PL relations for WLM in the J and K bands:

$$J = -3.153 \log P + (22.795 \pm 0.055) \sigma = 0.256$$

$$K = -3.261 \log P + (22.436 \pm 0.048) \sigma = 0.223$$

The dispersions are larger than the dispersions of the J and K band PL relations found in the LMC by Persson et al. (2004), which are 0.12 mag in both bands. These values should closely resemble the *intrinsic* dispersions of these relations, given that they are based on almost a hundred Cepheids whose mean magnitudes were derived from full infrared light curves. The dispersions of the present PL relations observed in WLM are larger mainly because they are based on a smaller number of stars, and on mean magnitudes which are less precise than in the LMC work of Persson et al. due to the existence of a few phase points only for each Cepheid. In order to determine the relative distance moduli between WLM and the LMC, we need to convert the NICMOS (LCO) photometric system used by Persson et al. (2004) to the UKIRT system utilized in this paper. According to Hawarden et al. (2001), there are just zero point offsets between the UKIRT and NICMOS systems (e.g. no color dependences) in the J and K filters, which amount to 0.034 and 0.015 mag, respectively. Applying these offsets, and assuming an LMC true distance modulus of 18.50 as in our previous work in the Araucaria Project, we derive distance moduli for WLM of 24.993 ± 0.055 mag in the J band, and 24.915 ± 0.045 mag in the K band.

As in our previous work, we will combine the distance moduli obtained from the near-infrared photometry with the values we had previously derived in the optical VI bands in Paper I, to obtain a very robust determination of both the true distance modulus of WLM, and the total (mean) reddening of the WLM Cepheids in our sample. When re-analyzing the results of Paper I in the course of this work, we detected that by accident we had interchanged

the distance modulus results reported for the V and Wesenheit bands (p. 601 of Paper I). The correct values for the reddened distance moduli in V and I derived from our photometry in Paper I are 25.156 ± 0.04 mag and 25.121 ± 0.03 mag, respectively. As in our previous papers in this series we adopt the extinction law of Schlegel et al. (1998) and fit a straight line to the relation $(m - M)_0 = (m - M)_\lambda - A_\lambda = (m - M)_\lambda - E(B - V) * R_\lambda$. Using the corrected, reddened distance moduli in the V and I photometric bands as given above, together with the values for the J and K bands obtained in this paper, we obtain the following values for the reddening and the true distance modulus of WLM from the multiwavelength analysis:

$$E(B - V) = 0.082 \pm 0.020$$

$$(m - M)_0 = 24.924 \pm 0.042,$$

This corresponds to a distance of WLM of 0.97 ± 0.02 Mpc.

In Table 4, we give the adopted values of R_λ and the unreddened, true distance moduli in each band which are obtained with the reddening value determined in our multi-wavelength solution. The agreement between the unreddened distance moduli obtained in each band is very good. In Fig. 6, we plot the apparent Cepheid distance moduli for WLM in VIJK as a function of R_λ , and the best fitting straight line to the data. It is appreciated that the total reddening, and the true distance modulus of WLM are indeed very well determined from this fit. By comparison with the foreground reddening of 0.02 mag (Schlegel et al. 1998) it is seen that most of the total average reddening of the WLM Cepheids is produced inside their host galaxy.

4. Discussion

As in the previous distance determinations of Local and Sculptor Group irregular and spiral galaxies from combined Cepheid near-infrared and optical photometry (see references cited in the Introduction) obtained in the Araucaria Project, our combined near-infrared and optical data for a sample of 31 Cepheids in WLM have led to a very robust distance determination for this relatively distant Local Group galaxy. Again, as for the other galaxies we have studied so far, we find that the slopes of the Cepheid PL relations in WLM in the near-infrared J and K bands are statistically in agreement with the one defined by the LMC Cepheids (Persson et al. 2004). Indeed, the free fits to the data discussed in section 3 of this paper yield slopes in both J and K which agree within 1σ with the adopted slope of Persson et al. (2004) from the LMC Cepheids. If we exclude the long-period Cepheid in the fits which carries a strong weight in the linear regression, the resulting values for the slopes

of the PL relation in J and K become -2.868 ± 0.426 , and -2.756 ± 0.368 , respectively. Due to their large uncertainties, which is a consequence of the now very restricted period range of the remaining 23 Cepheid sample, these values still agree within 1.5σ with the slopes of the Cepheid PL relation as determined in the LMC by Persson et al. (2004). The observed slopes of the IR PL relations in WLM are therefore consistent with the claim made by Gieren et al. (2005a) and others that the slope of the Cepheid PL relation is metallicity-independent, at least in near-infrared bands and for metallicities in the range from -0.35 dex appropriate for the LMC Cepheids (Luck et al. 1998) down to -0.8 to -1.0 dex, which is the metallicity of the young stellar population in WLM (Urbaneja et al. 2008), IC 1613 (Bresolin et al. 2007) and NGC 3109 (Evans et al. 2007) as determined from quantitative spectroscopy of blue supergiants. These results from our project support our adopted procedure to use the extremely well-determined slopes of the LMC Cepheid PL relations for the distance determination of our target galaxies.

The distance of WLM derived in this paper is 0.09 mag shorter than the one derived in Paper I from the Wesenheit index and agrees more closely with the distance for this galaxy derived from the I-band magnitude of the tip of the red giant branch. Indeed, Lee et al. (1993) obtained 24.87 ± 0.08 mag for the distance of WLM from the TRGB method, while Minniti & Zijlstra measured 24.75 ± 0.1 from their data. More recent results from the same method are 24.85 ± 0.08 mag obtained by McConnachie et al. (2005), and 24.93 ± 0.04 mag, obtained by Rizzi et al. (2007). Our own TRGB distance modulus of WLM reported in Paper I is 24.91 ± 0.08 . All these values are in excellent agreement with the WLM distance modulus of 24.92 obtained in this paper from the Cepheid VIJK photometry. Our Cepheid distance modulus is also in excellent agreement with the WLM distance modulus of 24.99 ± 0.10 mag derived very recently by Urbaneja et al. (2008) from a quantitative analysis of low resolution spectra of A and B supergiants via the flux-weighted gravity-luminosity relation of Kudritzki et al. (2003, 2008). Other distance measurements reported for WLM include the work of Valcheva et al. (2007) who obtained 24.84 ± 0.14 mag from IR photometry of 4 Cepheids, Rejkuba et al. (2000) who measured 24.95 ± 0.13 mag from the V magnitude of the horizontal branch in WLM, and Dolphin (2000) who obtained 24.88 ± 0.09 mag from fitting the V, I color-magnitude diagram of the galaxy. All these distance measurements agree within the uncertainties with the present Cepheid result.

In the previous Cepheid studies of our project in other Araucaria target galaxies the agreement between the distance obtained from the VI Wesenheit index, and the full multi-wavelength solution including the near-infrared bands was usually better (in the order of 2 %). Probably our result for WLM just demonstrates that a truly reliable and robust distance determination from Cepheids definitively requires the use of near-infrared photometry, and that the result from the (in principle) reddening-free VI Wesenheit magnitude might yield a

result which is still biased to some extent, particularly when blending with unresolved stars in the optical images becomes a problem. Once we have finished our VIJK distance analyses for all the target galaxies in our project, we will analyze this problem more exhaustively.

The distance modulus of WLM derived in this paper is not affected by any significant amount by the choice of the cutoff period adopted in the PL diagrams in Figs. 4 and 5. The zero points in the WLM PL relations in both J and K change by less than 0.03 mag if we use a longer period cutoff of $\log P$ (days) = 0.7 (leaving the 14 longest-period stars in the sample), or if we use the full sample of 30 Cepheids (excluding the one strongly blended Cepheid cep038). Specifically, retaining the full Cepheid sample yields a distance modulus of 24.902 ± 0.042 mag, whereas the sample of the 14 longest-period stars yields 24.947 ± 0.042 mag. Both values agree within their statistical 1σ errors with our adopted best distance modulus of 24.924 ± 0.042 mag. This probably demonstrates that our WLM Cepheid sample is large enough to fill the Cepheid instability strip in the HRD rather homogeneously, and that there are no overtone Cepheids among the shortest period variables in the sample, a conclusion which is supported by the asymmetric light curve shapes in V of all these variables.

We estimate that the combined effect of the different sources which contribute to the systematic uncertainty of our present multiwavelength Cepheid distance result for WLM generate an uncertainty not exceeding ± 0.06 mag, or 3%. These factors include the accuracy of the photometric zero points, the effect of blending on the Cepheid photometry, the effect of errors in the adopted reddening and a possible metallicity dependence of the Cepheid PL relation. The typical impact of these factors on the distance result has been discussed quite exhaustively in the previous papers of this series. Here we only want to stress that the zero point of the near-infrared photometry for WLM reported in this paper is probably even more accurate (0.01 mag) than for the other, previously studied target galaxies of our project, as a consequence of having data and independent photometric calibrations from 6 different photometric nights, which has been an especially fortunate circumstance. The impact of blending of the Cepheids on the distance was found to be less than 2% in the case of NGC 300 (Bresolin et al. 2005) which is at twice the distance of WLM, which allows us to conclude that its effect in the present case of WLM is certainly not larger than 2%, or 0.04 mag, and very likely smaller than this. We adopt 0.03 mag for this source of systematic error. Regarding the effect of reddening, our analysis and the results presented in Fig. 6 and Table 4 convincingly demonstrate that the total reddening suffered by the WLM Cepheids has been very accurately determined in this study, and any residual effect of reddening on our distance result is clearly less than 2%, or 0.04 mag. We want to stress again that this elimination of reddening as a very serious source of systematic error in Cepheid distances to galaxies based on optical photometry alone is probably the single most important advantage offered by our combined optical-infrared procedure to measure Cepheid distances to late-

type galaxies in our project. The influence of metallicity effects on our distance result is harder to estimate, at the present time. While we have presented evidence that the effect of metallicity differences on the slope of the PL relation is probably negligible, its effect on the zero points of the PL relation is not well established yet. We will investigate this question in detail in a forthcoming paper in which we will compare the distance results for all our target galaxies from the different stellar methods. However, previous work, like the one of Storm et al. (2004), seem to indicate that the metallicity effect on the PL relation zero point is quite small. We estimate its contribution to the systematic error of our WLM distance modulus to be 0.04 mag. Summing up the different contributions, we arrive at an estimated total ± 0.06 mag systematic uncertainty of our present WLM distance modulus.

As in the previous papers of this series, our distance result for WLM is tied to an *assumed* LMC distance modulus of 18.50 mag, which probably constitutes the single most important source of systematic uncertainty (e.g. Benedict et al. 2002, Schaefer 2008). If future work changes this adopted value of the LMC distance, we can easily adapt the distances of the target galaxies of our project to the new value. The *relative* distances between our different target galaxies will remain unaffected.

Finally, as remarked already before, our analyses of a number of galaxies with large differences in the metallicities of their young stellar populations keep indicating us so far that the slopes of the Cepheid PL relations in the various optical and particularly in the near-infrared J and K bands are identical and thus not dependent on metallicity. Subtle effects like the possible nonlinearity of the PL relation at 10 days as reported by Ngeow et al. (2008) do not seem to have any appreciable effect on the distance measurement with this tool, particularly in the near-infrared domain. The possible dependence of the zero points of the PL relation in different bands on metallicity will be studied in detail in a forthcoming paper of this series by comparison with the results from other distance indicators.

5. Conclusions

The main conclusions of this paper can be summarized as follows:

1. We have obtained deep near-infrared photometry of a sizeable sample (31 stars) of Cepheid variables in the Local Group galaxy WLM which were previously discovered in a wide-field imaging survey conducted in the optical V and I bands.
2. From our calibrated infrared photometry in J and K we derive tight period-luminosity relations for the Cepheids in WLM which are very well fitted by the slopes of the LMC Cepheid PL relations in these bands of Persson et al. (2004), supporting the conclusion

that for a broad range of metallicities the Cepheid PL relation in near-infrared bands is independent of metallicity.

3. Combining the reddened distance moduli in the J and K bands obtained in this paper with those obtained in Paper I in the V and I bands we derive a total reddening of the WLM Cepheids of $E(B-V) = 0.082 \pm 0.02$, and a absorption corrected, true distance modulus for WLM of 24.92 ± 0.04 mag (random error) ± 0.06 mag (systematic error).

4. We report on an error in Paper I where we had confused the distance modulus in the V band with the one in the Wesenheit band. For our present distance determination, we have obviously used the correct value of the reddened distance modulus in V as obtained from the data in Paper I. The true distance modulus from the Wesenheit magnitude obtained in that paper must read 25.014 ± 0.036 mag, rather than the erroneously given value of 25.144 mag.

5. The WLM distance derived from the multiwavelength VIJK Cepheid analysis in this paper is in excellent agreement with the various determinations of the WLM distance from the I-band tip of the red giant branch method and from other techniques which have been reported in the literature.

6. With the completion of the Cepheid multiwavelength distance determination for WLM there are now six late-type galaxies in the Araucaria Project for which such distance determinations have been carried out (WLM, IC 1613, NGC 3109 and NGC 6822 in the Local Group and NGC 300 and NGC 55 in the Sculptor Group; see references in the Introduction). Corresponding work for two more Sculptor Group galaxies, NGC 247 and NGC 7793, is underway. A comparison of these distances with those derived for the same galaxies from the other techniques we are using in our project will be conducted in due time and is expected to lead to an improved determination of the metallicity dependence of the different techniques. This will finally yield a set of very accurate distances to a number of nearby galaxies in the 0.05-4 Mpc range which can be used to calibrate more accurately other techniques which reach out to distances large enough for an improved determination of the Hubble constant.

WG, GP and DM gratefully acknowledge financial support for this work from the Chilean Center for Astrophysics FONDAF 15010003, and from the BASAL Centro de Astrofísica y Tecnologías Afines (CATA). Support from the Polish grant N203 002 31/046 and the FOCUS subsidy of the Foundation for Polish Science (FNP) is also acknowledged. It is a great pleasure to thank the support astronomers at ESO-La Silla and at Las Campanas Observatory for their expert help in the observations. We also thank the ESO OPC and CNTAC for allotting generous amounts of observing time to this project. Helpful comments of a referee on a previous version of this paper are appreciated.

REFERENCES

- Benedict, G.F., McArthur, B.E., Fredrick, L.W., et al., 2002, *AJ*, 123, 473
- Bono, G., Caputo, F., Castellani, V., and Marconi, M., 1999, *ApJ*, 512, 711
- Bresolin, F., Pietrzyński, G., Gieren, W., and Kudritzki, R.-P., 2005, *ApJ*, 634, 1020
- Bresolin, F., Pietrzyński, G., Urbaneja, M.A., Gieren, W., Kudritzki, R.P., and Venn, K., 2006, *ApJ*, 648, 1007
- Bresolin, F., Urbaneja, M.A., Gieren, W., Pietrzyński, G., and Kudritzki, R.P., 2007, *ApJ*, 671, 2028
- Dolphin, A.E., 2000, *ApJ*, 531, 804
- Evans, C.J., Bresolin, F., Urbaneja, M.A., Pietrzyński, G., Gieren, W., and Kudritzki, R.P., 2007, *ApJ*, 659, 1198
- Fouqué, P., Arriagada, P., Storm, J., Barnes, T.G., Nardetto, N., Merand, A., Kervella, P., Gieren, W., Bersier, D., Benedict, G.F., and McArthur, B.E., 2007, *A&A*, 476, 73
- Gieren, W., Storm, J., Barnes III, T.G., Fouqué, P., Pietrzyński, G., and Kienzle, F., 2005a, *ApJ*, 627, 224
- Gieren, W., Pietrzyński, G., Soszyński, I., Bresolin, F., Kudritzki, R.-P., Minniti, D., and Storm, J., 2005b, *ApJ*, 628, 695
- Gieren, W., Pietrzyński, G., Nalewajko, K., et al., 2006, *ApJ*, 647, 1056
- Gieren, W., Pietrzyński, G., Soszyński, I., Bresolin, F., Kudritzki, R.P., Storm, J., and Minniti, D., 2008, *ApJ*, 672, 266
- Hawarden, T.G., Leggett, S.K., Letawsky, M.B., et al., 2001, *MNRAS*, 325, 563
- Kudritzki, R.P., Bresolin, F., and Przybilla, N., 2003, *ApJ*, 582, L83
- Kudritzki, R.P., Urbaneja, M.A., Bresolin, F., Przybilla, N., Gieren, W., and Pietrzyński, G., 2008, *ApJ*, in press
- Lee, M.G., Freedman, W.L., and Madore, B.F., 1993, *ApJ*, 417, 553
- Luck, E., Moffett, T.J., Barnes, T.G., and Gieren, W., 1998, *AJ*, 115, 605
- Madore, B.F., and Freedman, W.L., 1991, *PASP*, 103, 933
- McConnachie, A.W., Irwin, M.J., Ferguson, A.M.N., Ibata, R.A., Lewis, G.F., and Tanvir, N., 2005, *MNRAS*, 356, 979
- McGonegal, R., McAlary, C.W., Madore, B.F., and McLaren, R.A., 1982, *ApJ*, 257, L33
- Minniti, D., and Zijlstra, A.A., 1997, *AJ*, 114, 147

- Ngeow, C.C., Kanbur, S.M., and Nanthakumar, A., 2008, *A&A*, 477, 621
- Persson, S.E., Madore, B.F., Krzeminski, W., Freedman, W.L., Roth, M, and Murphy, D.C., 2004, *AJ*, 128, 2239
- Pietrzyński, G., and Gieren, W., 2002, *AJ*, 124, 2633
- Pietrzyński, G., Gieren, W., Soszyński, I., Bresolin, F., Kudritzki, R.-P., Dall’Ora, M., Storm, J., and Bono, G., 2006, *ApJ*, 642, 216
- Pietrzyński, G., Gieren, W., Udalski, A., et al., 2007, *AJ*, 134, 594 (Paper I)
- Rejkuba, M., Minniti, D., Gregg, M.D., Zijlstra, A.A., Alonso, M.V., and Goudfrooij, P., 2000, *AJ*, 120, 801
- Rizzi, L., Tully, R.B., Makarov, D., et al., 2007, *ApJ*, 661, 815
- Sandage, A., and Carlson, G., 1985, *AJ*, 90, 1464
- Schaefer, B.E., 2008, *AJ*, 135, 112
- Schlegel, D.J., Finkbeiner, D.P., and Davis, M., 1998, *ApJ*, 500, 525
- Soszyński, I., Gieren, W., and Pietrzyński, G., 2005, *PASP*, 117, 823
- Soszyński, I., Gieren, W., Pietrzyński, G., Bresolin, F., Kudritzki, R.P, and Storm, J., 2006, *ApJ*, 648, 375
- Storm, J., Carney, B.W., Gieren, W., Fouqué, P., Fry, A.M., and Latham, D.W., 2004, *A&A*, 415, 531
- Urbaneja, M.A., Kudritzki, R.P., Bresolin, F., Przybilla, N., Gieren, W. and Pietrzyński, G., 2008, *ApJ*, submitted
- Valcheva, A.T., Ivanov, V.D., Ovcharov, E.P., and Nedialkov, P., 2007, *A&A*, 466, 501




Fig. 1.— The location of the observed fields in WLM on the DSS blue plate. We observed one NTT/SOFI field, and three Magellan/PANIC fields (see text). The fields contain 31 Cepheid variables. The size of the SOFI field (S1) is 4.9×4.9 arcmin.




Fig. 2.— An example of the comparison of the photometric zero points in the K and J bands of WLM obtained with the NTT and SOFI on two different nights and calibrated completely independently. The difference in zero points for the data obtained for other nights with SOFI and PANIC are very similar and are always smaller than 0.02 mag, which reflects the very good quality of our calibrations.




Fig. 3.— Comparison of our photometry with the 2MASS data. In spite of the large scatter towards the fainter magnitudes caused by the low accuracy of the 2MASS photometry of faint stars, no clear evidence for significant offsets in the zero points is present.




Fig. 4.— Cepheid period-luminosity relations for WLM in the J band. The 24 variables with periods larger than 3.16 days (filled circles) have been adopted for the distance determination. The slope of the fit to a line has been adopted from the work of Persson et al. on the LMC Cepheids, and fits the WLM Cepheid data very well.




Fig. 5.— Same as Fig. 4, for the K band.




Fig. 6.— Apparent distance moduli to WLM as derived in different photometric bands, plotted against the ratio of total to selective extinction as adopted from the Schlegel et al. reddening law. The intersection and slope of the best-fitting line give the true distance modulus of WLM, and the total mean reddening of its Cepheids, respectively.

Table 1.

Field ID	Field name	RA 2000	DEC 2000	Instrument
S1	WLM-F1	00:01:57.67	-15:25:29.5	SOFI
P1	WLM-F1	00:01:58.26	-15:24:40.2	PANIC
P2	WLM-F2	00:01:54.98	-15:27:01.8	PANIC
P3	WLM-F3	00:01:54.87	-15:29:48.9	PANIC

Table 2. Journal of the Individual J and K band Observations of WLM

Star ID	Field ID	Period (days)	J MJD -2400000	J (mag)	σ (mag)	K MJD -2400000	K (mag)	σ (mag)
cep001	S1	54.17118	54001.08144853	17.309	0.024	54001.09783376	16.955	0.018
cep001	S1	54.17118	54002.10563506	17.327	0.019	54002.11759493	16.802	0.017
cep001	P1	54.17118	54427.06900000	17.341	0.004	54427.08676000	16.781	0.007
cep001	P1	54.17118	54428.14376000	17.326	0.004	54428.15841000	16.854	0.006
cep001	S1	54.17118	54429.01840384	17.313	0.014	54429.03095727	16.813	0.027
cep001	S1	54.17118	54430.01460106	17.292	0.013	54430.02712407	16.807	0.023
cep002	P3	10.34249	54428.06006000	19.737	0.020	54428.07505000	19.344	0.034
cep003	P2	10.33645	54427.15887000	19.677	0.025	54427.17380000	19.277	0.062
cep005	P3	8.63110	54428.06006000	19.916	0.020	54428.07505000	19.469	0.039
cep007	S1	8.12051	54001.08144853	19.927	0.083	54001.09783376	19.463	0.077
cep007	S1	8.12051	54002.10563506	19.984	0.063	54002.11759493	19.561	0.100
cep007	P1	8.12051	54427.06900000	20.182	0.019	54427.08676000	19.814	0.048
cep007	P1	8.12051	54428.14376000	20.126	0.022	54428.15841000	19.620	0.054
cep007	S1	8.12051	54429.01840384	19.840	0.078	54429.03095727	19.511	0.192
cep007	S1	8.12051	54430.01460106	20.001	0.104	54430.02712407	19.383	0.150
cep008	P2	7.49672	54427.15887000	20.040	0.029	54427.17380000	19.351	0.048
cep010	S1	7.32485	54001.08144853	20.167	0.092	54001.09783376	19.858	0.088
cep010	S1	7.32485	54002.10563506	19.868	0.050	54002.11759493	19.472	0.107
cep010	P1	7.32485	54427.06900000	19.955	0.024	54427.08676000	19.654	0.054
cep010	P1	7.32485	54428.14376000	19.899	0.030	54428.15841000	19.525	0.056
cep011	P3	6.64055	54428.06006000	20.187	0.018	54428.07505000	19.895	0.042
cep012	S1	6.15754	54001.08144853	19.841	0.083	54001.09783376	19.263	0.065
cep012	S1	6.15754	54002.10563506	19.930	0.071	54002.11759493	19.388	0.087
cep012	P1	6.15754	54427.06900000	99.999	9.999	54427.08676000	19.438	0.051
cep012	P1	6.15754	54428.14376000	20.131	0.040	54428.15841000	19.549	0.054
cep012	S1	6.15754	54429.01840384	19.770	0.075	54429.03095727	19.141	0.147
cep013	P3	6.05309	54428.06006000	20.339	0.027	54428.07505000	19.989	0.062

Table 2—Continued

Star ID	Field ID	Period (days)	J MJD -2400000	J (mag)	σ (mag)	K MJD -2400000	K (mag)	σ (mag)
cep015	S1	5.43153	54001.08144853	20.626	0.124	54001.09783376	20.284	0.138
cep015	S1	5.43153	54002.10563506	20.638	0.091	54002.11759493	20.322	0.190
cep015	P1	5.43153	54427.06900000	20.574	0.028	54427.08676000	20.320	0.093
cep015	P1	5.43153	54428.14376000	20.677	0.036	54428.15841000	20.280	0.087
cep015	S1	5.43153	54430.01460106	20.386	0.131	54430.02712407	19.793	0.209
cep016	S1	5.20796	54001.08144853	20.664	0.120	54001.09783376	20.099	0.102
cep016	S1	5.20796	54002.10563506	20.582	0.071	54002.11759493	20.016	0.131
cep016	P1	5.20796	54427.06900000	20.385	0.022	54427.08676000	20.090	0.078
cep016	P1	5.20796	54428.14376000	20.493	0.035	54428.15841000	19.980	0.071
cep016	S1	5.20796	54430.01460106	20.404	0.134	54430.02712407	20.163	0.274
cep017	P3	5.12851	54428.06006000	20.727	0.025	54428.07505000	20.332	0.060
cep018	S1	5.02134	54001.08144853	20.555	0.118	54001.09783376	20.402	0.140
cep018	S1	5.02134	54002.10563506	20.567	0.080	54002.11759493	20.328	0.173
cep018	S1	5.02134	54429.01840384	20.555	0.149	54429.03095727	19.730	0.226
cep019	S1	4.92341	54001.08144853	20.338	0.107	54001.09783376	20.217	0.138
cep019	S1	4.92341	54002.10563506	20.494	0.086	54002.11759493	19.990	0.141
cep019	P1	4.92341	54427.06900000	20.379	0.031	54427.08676000	19.964	0.070
cep019	P1	4.92341	54428.14376000	20.626	0.037	54428.15841000	20.379	0.098
cep019	S1	4.92341	54429.01840384	20.265	0.113	54429.03095727	19.828	0.260
cep020	S1	4.91559	54429.01840384	99.999	9.999	54429.03095727	20.413	0.409
cep021	S1	4.86831	54002.10563506	21.096	0.115	54002.11759493	20.345	0.186
cep022	S1	4.71140	54001.08144853	20.726	0.136	54001.09783376	20.232	0.141
cep022	S1	4.71140	54002.10563506	20.735	0.097	54002.11759493	20.234	0.160
cep022	P1	4.71140	54427.06900000	20.782	0.040	54427.08676000	20.301	0.079
cep022	P1	4.71140	54428.14376000	20.511	0.032	54428.15841000	20.492	0.098
cep023	S1	4.61967	54001.08144853	20.597	0.122	54001.09783376	20.607	0.171
cep023	S1	4.61967	54002.10563506	20.669	0.093	54002.11759493	20.222	0.177

Table 2—Continued

Star ID	Field ID	Period (days)	J MJD -2400000	J (mag)	σ (mag)	K MJD -2400000	K (mag)	σ (mag)
cep023	P1	4.61967	54427.06900000	20.630	0.026	54427.08676000	20.316	0.085
cep023	P1	4.61967	54428.14376000	20.646	0.039	54428.15841000	20.328	0.086
cep024	P2	4.36958	54427.15887000	20.594	0.048	54427.17380000	20.307	0.148
cep024	S1	4.36958	54430.01460106	99.999	9.999	54430.02712407	19.658	0.174
cep026	S1	3.97606	54001.08144853	21.726	0.299	54001.09783376	21.334	0.333
cep026	S1	3.97606	54002.10563506	21.617	0.180	54002.11759493	21.026	0.307
cep026	S1	3.97606	54429.01840384	99.999	9.999	54429.03095727	20.454	0.468
cep026	S1	3.97606	54430.01460106	99.999	9.999	54430.02712407	20.631	0.387
cep027	P2	3.86503	54427.15887000	20.770	0.048	54427.17380000	20.654	0.178
cep027	S1	3.86503	54429.01840384	99.999	9.999	54429.03095727	20.116	0.335
cep031	S1	3.65377	54001.08144853	20.290	0.092	54001.09783376	20.067	0.098
cep031	S1	3.65377	54002.10563506	20.487	0.064	54002.11759493	20.049	0.126
cep031	P1	3.65377	54427.06900000	20.401	0.029	54427.08676000	19.914	0.072
cep031	P1	3.65377	54428.14376000	20.533	0.045	54428.15841000	20.201	0.102
cep031	S1	3.65377	54429.01840384	20.684	0.154	54429.03095727	20.014	0.290
cep031	S1	3.65377	54430.01460106	20.169	0.106	54430.02712407	19.756	0.181
cep033	S1	3.30475	54001.08144853	99.999	9.999	54001.09783376	20.533	0.155
cep033	P1	3.30475	54428.14376000	21.006	0.074	54428.15841000	20.300	0.112
cep033	S1	3.30475	54430.01460106	99.999	9.999	54430.02712407	20.861	0.512
cep036	S1	3.13908	54001.08144853	21.178	0.182	54001.09783376	20.372	0.136
cep036	P1	3.13908	54427.06900000	21.180	0.051	54427.08676000	21.031	0.166
cep037	S1	3.13102	54001.08144853	21.301	0.205	54001.09783376	20.643	0.169
cep037	P1	3.13102	54427.06900000	21.067	0.040	54427.08676000	20.774	0.123
cep037	P1	3.13102	54428.14376000	21.264	0.058	54428.15841000	20.769	0.136
cep037	S1	3.13102	54429.01840384	99.999	9.999	54429.03095727	20.195	0.357
cep038	P1	3.03891	54427.06900000	19.839	0.018	54427.08676000	18.954	0.029
cep038	P1	3.03891	54428.14376000	19.834	0.019	54428.15841000	19.022	0.028

Table 2—Continued

Star ID	Field ID	Period (days)	J MJD -2400000	J (mag)	σ (mag)	K MJD -2400000	K (mag)	σ (mag)
cep038	S1	3.03891	54429.01840384	19.957	0.084	54429.03095727	19.021	0.132
cep038	S1	3.03891	54430.01460106	19.899	0.093	54430.02712407	19.091	0.118
cep039	S1	3.02576	54001.08144853	21.155	0.184	54001.09783376	21.439	0.374
cep039	P1	3.02576	54427.06900000	21.061	0.044	54427.08676000	20.562	0.164
cep039	S1	3.02576	54429.01840384	21.796	0.431	54429.03095727	20.483	0.428
cep041	P3	2.92899	54428.06006000	21.333	0.072	54428.07505000	21.107	0.154
cep043	P3	2.88896	54428.06006000	20.906	0.051	54428.07505000	20.723	0.130
cep049	S1	2.51004	54001.08144853	21.440	0.235	54001.09783376	21.382	0.377
cep049	P1	2.51004	54427.06900000	20.947	0.037	54427.08676000	20.796	0.133
cep049	P1	2.51004	54428.14376000	21.108	0.052	54428.15841000	20.838	0.127

Table 3. Intensity mean J and K magnitudes for 30 Cepheid variables in WLM

Star ID	Period (days)	J (mag)	σ (mag)	K (mag)	σ (mag)
cep001	54.17118	17.318	0.015	16.835	0.018
cep002	10.34249	19.737	0.020	19.344	0.034
cep003	10.33645	19.677	0.025	19.277	0.062
cep005	8.63110	19.916	0.020	19.469	0.039
cep007	8.12051	20.010	0.069	19.559	0.116
cep008	7.49672	20.040	0.029	19.351	0.048
cep010	7.32485	19.972	0.056	19.627	0.079
cep011	6.64055	20.187	0.018	19.895	0.042
cep012	6.15754	19.918	0.062	19.356	0.088
cep013	6.05309	20.339	0.027	19.989	0.062
cep015	5.43153	20.580	0.093	20.200	0.152
cep016	5.20796	20.506	0.088	20.070	0.151
cep017	5.12851	20.727	0.025	20.332	0.060
cep018	5.02134	20.559	0.119	20.153	0.183
cep019	4.92341	20.420	0.082	20.076	0.156
cep020	4.91559	99.999	9.999	20.413	0.409
cep021	4.86831	21.096	0.115	20.345	0.186
cep022	4.71140	20.689	0.087	20.315	0.124
cep023	4.61967	20.636	0.080	20.368	0.137
cep024	4.36958	20.594	0.034	19.983	0.162
cep026	3.97606	21.672	0.175	20.861	0.379
cep027	3.86503	20.770	0.034	20.385	0.268
cep031	3.65377	20.427	0.092	20.000	0.162
cep033	3.30475	21.006	0.043	20.565	0.316
cep036	3.13908	21.179	0.134	20.702	0.152
cep037	3.13102	21.211	0.108	20.595	0.218
cep038	3.03891	19.882	0.064	19.022	0.091
cep039	3.02576	21.337	0.272	20.828	0.342
cep041	2.92899	21.333	0.072	21.107	0.154
cep043	2.88896	20.906	0.051	20.723	0.130

Table 3—Continued

Star ID	Period (days)	J (mag)	σ (mag)	K (mag)	σ (mag)
cep049	2.51004	21.165	0.141	21.005	0.242

Table 4. Reddened and Absorption-Corrected Distance Moduli for WLM in Optical and Near-Infrared Bands

Band	V	I	J	K	$E(B - V)$
$m - M$	25.156	25.121	24.993	24.915	–
R_λ	3.24	1.96	0.902	0.367	–
$(m - M)_0$	24.896	24.964	24.920	24.885	0.08